



Soybean Frogeye Leaf Spot Risk Forecast

For period of Aug 8 – Sept 6, Date of Issue: Aug 9, 2020

One-Month Weather Outlook

Over the next 30 days both temperatures and precipitation will be normal or above normal for the entire South except for Florida. During the next two weeks temperatures will generally be normal with above normal precipitation except for Florida.

Frogeye Leaf Spot Risk Summary

The risk of frogeye leaf spot is generally high and at similar risk levels to the previous forecast for late-planted soybeans




Next two-week risk interpretation

During the next two weeks the FLS risk is moderate to high for regions west of the Appalachian Mountains. In areas of orange and extending into areas of yellow, consider treatment of susceptible cultivars that were planted late.

ONE-MONTH RISK MAP

WHAT TO DO?	
HIGH	Consider treatment, especially susceptible cultivars
MODERATE	Scout fields and weigh disease risk factors
LOW OR VERY LOW	Monitor risk forecasts



DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Stratego® YLD are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.

Risk Level



Bayer CropScience



STRATEGO® YLD



Asian Soybean Rust Risk Forecast

For period of Aug 8 – Sept 6, Date of Issue: Aug 9, 2020

One-Month Weather Outlook

Over the next 30 days both temperatures and precipitation will be normal or above normal for the entire South except for Florida. During the next two weeks temperatures will generally be normal with above normal precipitation except for Florida.

Asian Soybean Rust Risk Summary

The risk of Asian soybean rust in the southern states is low for most areas.


Next two-week risk interpretation

The risk of soybean rust is low in the south except for some spots having weather conditions moderately favorable to the disease. In areas of yellow, scout late-planted soybeans.



ONE-MONTH RISK MAP

WHAT TO DO?	
HIGH	Consider treatment if SBR is confirmed in the area
MODERATE	Scout fields and weigh disease risk factors
LOW OR VERY LOW	Monitor risk forecasts



DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Stratego® YLD are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.

Risk Level



Bayer CropScience



Corn Southern Rust Risk Forecast

For period of Aug 8 – Sept 6, Date of Issue: Aug 9, 2020

One-Month Weather Outlook


Over the next 30 days both temperatures and precipitation will be normal or above normal for the entire South except for Florida. During the next two weeks temperatures will generally be normal with above normal precipitation except for Florida.

Southern Rust Risk Summary

The risk of southern corn rust is moderate to high in the South.


Next 2-week risk interpretation

Conditions are very conducive to SCR development. If your crop has yet to reach R5 (dent), consider treatment if the disease is present. Do not anticipate that a fungicide application will eliminate existing infections, but it may limit the extent of the yield damage.



ONE-MONTH RISK MAP

WHAT TO DO?	
HIGH	Consider treatment if CSR is confirmed in the area
MODERATE	Scout fields and weigh disease risk factors
LOW OR VERY LOW	Monitor risk forecasts



Risk Level



DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Delaro® are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.

