



DELARO[®]
Complete
 458 SC Fungicide

Soybean White Mold Risk Forecast

For period of June 12 – July 10, Date of Issue: June 13, 2021

One-Month Weather Outlook

Over the next 30 days temperatures will be in normal ranges along with normal precipitation for most of the Corn Belt. During the next two weeks temperatures will be near normal except for northern MN and ND where it will be cooler than average. Above average precipitation is predicted in most sections of the Corn Belt.

White Mold Risk Summary


The risk of soybean white mold is low except for the far northern counties of the forecast map.



Next two-week risk interpretation

The risk of soybean white mold is low in the northern soybean production regions for the next two weeks. In the northernmost counties the conditions are moderately favorable for disease development on early-planted soybeans.

ONE-MONTH RISK MAP



WHAT TO DO?

HIGH	Treat flowering fields that have a closing canopy and a history of white mold infections
MODERATE	Scout fields and weigh disease risk factors
LOW OR VERY LOW	Monitor risk forecasts

DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Delaro[®] are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.

Risk Level



RESTRICTED



Soybean Brown Spot Risk Forecast

For period of June 12 – July 10, Date of Issue: June 13, 2021

One-Month Weather Outlook

Over the next 30 days temperatures will be in normal ranges along with normal precipitation for most of the Corn Belt. During the next two weeks temperatures will be near normal except for northern MN and ND where it will be cooler than average. Above average precipitation is predicted in most sections of the Corn Belt.

Brown Spot Risk Summary

The risk of soybean brown spot is low to moderate in the northern soybean production regions.


Next two-week risk interpretation

For the next two weeks the risk of soybean brown spot is moderate in some areas of the Corn Belt. In region of yellow, disease may be found on the lower leaves of the canopy in early-planted soybean fields. Scout fields in yellow, especially fields which have had severe brown spot previously.



ONE-MONTH RISK MAP

WHAT TO DO?	
HIGH	Consider treatment, especially susceptible cultivars
MODERATE	Scout fields and weigh disease risk factors
LOW OR VERY LOW	Monitor risk forecasts



Risk Level

Very-low	Low	Moderate	High
----------	-----	----------	------

DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Delaro® are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.



RESTRICTED



Soybean Frogeye Leaf Spot Risk Forecast

For period of June 12 – July 10, Date of Issue: June 13, 2021

One-Month Weather Outlook

Over the next 30 days temperatures will be in normal ranges along with normal precipitation for most of the Corn Belt. During the next two weeks temperatures will be near normal except for northern MN and ND where it will be cooler than average. Above average precipitation is predicted in most sections of the Corn Belt.

Frogeye Leaf Spot Risk Summary

The risk of soybean frogeye leaf spot is low in the Corn Belt during this period.



Next two-week risk interpretation

For the next two weeks the risk of FLS is low in the Corn Belt as weather conditions are not favorable to the development of this disease. Monitor our forecasts as soybeans progress.

ONE-MONTH RISK MAP



WHAT TO DO?

HIGH	Consider treatment, especially susceptible cultivars
MODERATE	Scout fields and weigh disease risk factors
LOW OR VERY LOW	Monitor risk forecasts

DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Delaro® are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.

Risk Level

