



DELARO®
Complete
458 SC Fungicide

Soybean White Mold Risk Forecast

For period of June 19 – July 17, Date of Issue: June 20, 2021

One-Month Weather Outlook


Over the next 30 days temperatures will be generally normal south of I-80 and warmer to the north. Precipitation will be in the normal range for the Corn Belt. During the next two weeks temperatures will be cooler than normal with higher than normal rainfall east of the Missouri River except for the western half of Minnesota.

White Mold Risk Summary


The risk of soybean white mold remains low except for some northernmost areas.

Next two-week risk interpretation

The risk of soybean white mold is low for soybean production during the next two weeks. In some farther north areas (yellow), conditions are moderately favorable for infections, especially for fields planted early. In areas of yellow, scout early-planted fields for white mold mushrooms if soil moisture is good.



ONE-MONTH RISK MAP



WHAT TO DO?	
HIGH	Treat flowering fields that have a closing canopy and a history of white mold infections
MODERATE	Scout fields and weigh disease risk factors
LOW OR VERY LOW	Monitor risk forecasts

DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Delaro® are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.

Risk Level



RESTRICTED



Soybean Brown Spot Risk Forecast

For period of June 19 – July 17, Date of Issue: June 20, 2021

One-Month Weather Outlook

Over the next 30 days temperatures will be generally normal south of I-80 and warmer to the north. Precipitation will be in the normal range for the Corn Belt. During the next two weeks temperatures will be cooler than normal with higher than normal rainfall east of the Missouri River except for the western half of Minnesota.

Brown Spot Risk Summary

The risk of soybean brown spot is moderate for the southern and western areas of the forecast map.


Next two-week risk interpretation

For the next two weeks the risk of soybean brown spot is moderate in the southern half of the Corn Belt. In region of yellow, disease may be found in the lower canopy of early-planted soybeans. Scout fields in yellow, especially for fields which had severe brown spot previously.



ONE-MONTH RISK MAP

WHAT TO DO?	
HIGH	Consider treatment, especially susceptible cultivars
MODERATE	Scout fields and weigh disease risk factors
LOW OR VERY LOW	Monitor risk forecasts



Risk Level

Very-low	Low	Moderate	High
----------	-----	----------	------

DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Delaro® are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.



RESTRICTED



Soybean Frogeye Leaf Spot Risk Forecast

For period of June 19 – July 17, Date of Issue: June 20, 2021

One-Month Weather Outlook

Over the next 30 days temperatures will be generally normal south of I-80 and warmer to the north. Precipitation will be in the normal range for the Corn Belt. During the next two weeks temperatures will be cooler than normal with higher than normal rainfall east of the Missouri River except for the western half of Minnesota.

Frogeye Leaf Spot Risk Summary


The risk of soybean frogeye leaf spot is moderate in the southwestern corner of the forecast map.



Next two-week risk interpretation

During the next two weeks the risk of FLS is moderate in Kansas and western Missouri and low in the rest of the Corn Belt. In regions of yellow, scout susceptible cultivars and consider fungicide treatments if FLS is found, especially on fields with a history of FLS infections.

ONE-MONTH RISK MAP



WHAT TO DO?

HIGH	Consider treatment, especially susceptible cultivars
MODERATE	Scout fields and weigh disease risk factors
LOW OR VERY LOW	Monitor risk forecasts

DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Delaro® are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.

Risk Level

