



DELARO[®]
Complete
 458 SC Fungicide

Soybean White Mold Risk Forecast

For period of June 26 – July 24, Date of Issue: June 27, 2021

One-Month Weather Outlook


Over the next 30 days temperatures will be generally normal south of I-80 and warmer than normal for regions farther north. Precipitation will be in the normal range for the Corn Belt east of the Mississippi River. For the next two weeks temperatures are expected to be cooler than normal with higher than normal precipitations south of I-80, and warmer and dryer than normal north of I-80.

White Mold Risk Summary

The risk of soybean white mold remains low except for some northern areas.


Next two-week risk interpretation

The risk of soybean white mold is low to very low for much of the northern soybean production region during the next two weeks. In the far northern areas in yellow, conditions are moderately favorable to infections, especially for fields planted early in narrow rows.



ONE-MONTH RISK MAP

WHAT TO DO?	
HIGH	Treat flowering fields that have a closing canopy and a history of white mold infections
MODERATE	Scout fields and weigh disease risk factors
LOW OR VERY LOW	Monitor risk forecasts



Risk Level



DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Delaro[®] are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.



RESTRICTED



Soybean Brown Spot Risk Forecast

For period of June 26 – July 24, Date of Issue: June 27, 2021

One-Month Weather Outlook

Over the next 30 days temperatures are to be normal for regions south of I-80 and warmer than normal for regions north. Precipitations are to be in normal range for Corn Belt east of Mississippi. In next two weeks, temperatures are to be cooler than normal with higher than normal precipitations for regions south of I-80. Regions of north are warmer and dryer than normal.

Brown Spot Risk Summary

The risk of soybean brown spot is moderate for the Corn Belt with some high-risk spots.


Next two-week risk interpretation

For the next two weeks the risk of soybean brown spot is moderate in the Corn Belt. In areas of yellow, disease may be found in the lower canopy. Scout fields, especially fields with a history of severe brown spot, and consider spraying if the disease moves upward into the middle canopy.



ONE-MONTH RISK MAP

WHAT TO DO?	
HIGH	Consider treatment, especially susceptible cultivars
MODERATE	Scout fields and weigh disease risk factors
LOW OR VERY LOW	Monitor risk forecasts



Risk Level

Very-low	Low	Moderate	High
----------	-----	----------	------

DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Delaro® are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.



RESTRICTED



Soybean Frogeye Leaf Spot Risk Forecast

For period of June 26 – July 24, Date of Issue: June 27, 2021

One-Month Weather Outlook

Over the next 30 days temperatures are to be normal for regions south of I-80 and warmer than normal for regions north. Precipitations are to be in normal range for Corn Belt east of Mississippi. In next two weeks, temperatures are to be cooler than normal with higher than normal precipitations for regions south of I-80. Regions of north are warmer and dryer than normal.

Frogeye Leaf Spot Risk Summary

The risk of soybean frogeye leaf spot is moderate or high for the southcentral areas of the Corn Belt.


Next two-week risk interpretation

For the next two weeks the risk of FLS is high in parts of KS, MO and IL and moderate for a larger area of the Corn Belt. In areas of orange, consider spraying susceptible cultivars. In yellow areas scout susceptible cultivars and treat if FLS is found on the upper leaves, especially on fields that had FLS in prior seasons.



ONE-MONTH RISK MAP

WHAT TO DO?	
HIGH	Consider treatment, especially susceptible cultivars
MODERATE	Scout fields and weigh disease risk factors
LOW OR VERY LOW	Monitor risk forecasts



DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Delaro® are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.

Risk Level

