



Soybean Frogeye Leaf Spot Risk Forecast

For period of June 26 – July 24, Date of Issue: June 27, 2021

One-Month Weather Outlook

Over the next 30 days temperatures will hover near normal with higher than normal rainfall across the South. During the next two weeks temperatures will be cooler than average and precipitation will be above normal except for areas along the east coast where temperatures and precipitation will be normal.

Frogeye Leaf Spot Risk Summary

The high-risk areas for frogeye leaf spot continue to expand eastward across the South.


Next two-week risk interpretation

For the next two weeks the FLS risk is high for the western half of the southern states and moderate for the rest. In orange areas spray susceptible cultivars with a fungicide containing multiple modes of action. In region of yellow, scout susceptible cultivars and consider spraying if symptoms are observed on the top leaves.



ONE-MONTH RISK MAP

WHAT TO DO?	
HIGH	Consider treatment, especially susceptible cultivars
MODERATE	Scout fields and weigh disease risk factors
LOW OR VERY LOW	Monitor risk forecasts



DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Stratego® YLD are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.

Risk Level





STRATEGO[®] YLD



Asian Soybean Rust Risk Forecast

For period of June 26 – July 24, Date of Issue: June 27, 2021

One-Month Weather Outlook

Over the next 30 days temperatures will hover near normal with higher than normal rainfall across the South. During the next two weeks temperatures will be cooler than average and precipitation will be above normal except for areas along the east coast where temperatures and precipitation will be normal.

Asian Soybean Rust Risk Summary

The risk of soybean rust remains moderate for much of the southern states, with pockets at high risk.

Next two-week risk interpretation


For the next two weeks the weather conditions are favorable to SBR infections for the southern states east of the Mississippi River. In regions of yellow, spore loads are low, leading to low levels of infection. There is uncertainty whether these infections can become established and lead to late summer development. In areas of orange, spray if soybean rust is found.



ONE-MONTH RISK MAP

WHAT TO DO?

HIGH	Consider treatment if SBR is confirmed in the area
MODERATE	Scout fields and weigh disease risk factors
LOW OR VERY LOW	Monitor risk forecasts



DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Stratego[®] YLD are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.

Risk Level



Bayer CropScience



DELARO[®]
Complete
458 SC Fungicide

Corn Southern Rust Risk Forecast

For period of June 26 – July 24, Date of Issue: June 27, 2021

One-Month Weather Outlook


Over the next 30 days temperatures will hover near normal with higher than normal rainfall across the South. During the next two weeks temperatures will be cooler than average and precipitation will be above normal except for areas along the east coast where temperatures and precipitation will be normal.

Southern Rust Risk Summary

The risk of southern corn rust in the coastal areas has increased for this period.

Next 2-week risk interpretation


For the next two weeks the risk of southern corn rust is high for the coastal areas and generally moderate for the rest of the forecast region as weather conditions become favorable for the development of the disease. In regions of orange, treat fields where the disease has been observed. In regions of yellow, scout the fields and consider spraying if SCR is found or reported nearby.



ONE-MONTH RISK MAP

WHAT TO DO?

HIGH	Consider treatment if CSR is confirmed in the area
MODERATE	Scout fields and weigh disease risk factors
LOW OR VERY LOW	Monitor risk forecasts



Risk Level



DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Delaro[®] are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.



RESTRICTED



DELARO[®]
Complete
458 SC Fungicide

Northern Corn Leaf Blight Risk Forecast

For period of June 26 – July 24, Date of Issue: June 27, 2021

One-Month Weather Outlook

Over the next 30 days temperatures will hover near normal with higher than normal rainfall across the South. During the next two weeks temperatures will be cooler than average and precipitation will be above normal except for areas along the east coast where temperatures and precipitation will be normal.

Northern Leaf Blight Risk Summary

The risk of NCLB is at a similar level to the previous forecast, moderate to high.

Next two-week risk interpretation


The risk is moderate to high in most of the southern states. In regions of orange, spray susceptible hybrids if NCLB is detected or if the fields have a history of the disease. In regions of yellow, scout susceptible hybrids and consider spraying if NCLB is observed.



ONE-MONTH RISK MAP

WHAT TO DO?

HIGH	Treat susceptible hybrids and weigh risk factors for other fields
MODERATE	Scout fields and treat as conditions warrant
LOW OR VERY LOW	Monitor risk forecasts



Risk Level



DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Delaro[®] are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.



RESTRICTED



Corn Southern Rust Risk Forecast

For period of June 26 – July 24, Date of Issue: June 27, 2021

One-Month Weather Outlook


Over the next 30 days temperatures will be normal to cooler than normal, and rainfall will be normal to above normal across the state of Texas. For the next two weeks temperatures will be below average with above average precipitation.

Southern Rust Risk Summary


The risk of southern corn rust remains moderate to high for the entire state.

Next 2-week risk interpretation

The risk of southern corn rust in Texas remains moderate to high as conditions favor the development of this disease. In areas of orange, consider treatment. In areas of yellow, scout for the disease and consider spraying if the disease is found or reported nearby.



ONE-MONTH RISK MAP



WHAT TO DO?	
HIGH	Consider treatment if CSR is confirmed in the area
MODERATE	Scout fields and weigh disease risk factors
LOW OR VERY LOW	Monitor risk forecasts

DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. . **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Delaro® are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.

Risk Level



RESTRICTED



DELARO[®]
Complete
458 SC Fungicide

Northern Corn Leaf Blight Risk Forecast

For period of June 26 – July 24, Date of Issue: June 27, 2021

One-Month Weather Outlook

Over the next 30 days temperatures will be normal to cooler than normal, and rainfall will be normal to above normal across the state of Texas. For the next two weeks temperatures will be below average with above average precipitation.

Northern Leaf Blight Risk Summary


The risk of northern corn leaf blight is moderate in eastern Texas and low for the rest of the state.

Next two-week risk interpretation

Weather conditions in eastern Texas are moderately favorable for NCLB and unfavorable in the west as temperatures increase. In regions of yellow, scout susceptible hybrids and consider treating if NCLB incidence is rising or the fields have a history of disease infections.



ONE-MONTH RISK MAP



Risk Level



WHAT TO DO?

HIGH	Treat susceptible hybrids and weigh risk factors for other fields
MODERATE	Scout fields and treat as conditions warrant
LOW OR VERY LOW	Monitor risk forecasts

DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Delaro[®] are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.



RESTRICTED