



# DELARO Complete 458 SC Fungicide

# **Soybean White Mold Risk Forecast**

For period of July 10 - Aug 8, Date of Issue: July 11, 2021

#### **One-Month Weather Outlook**


The next 30 days will be warmer and drier than normal for the northern parts of the Corn Belt, and generally normal for the southern parts. During the next two weeks temperatures will be cooler than normal with higher than normal precipitation for areas south of I-80. North of I-80 it will be warmer and dryer than normal except for the northeastern areas.

# **White Mold Risk Summary**

The risk of soybean white mold is moderate around the Great Lakes and in northern MN.


## **Next two-week risk interpretation**

The conditions are moderately favorable for white mold in areas along the Great Lake and extending into northern Minnesota for the next two weeks. In region of yellow, scout for white mold mushrooms and treat before canopy closure. The risk is higher than moderate for fields with narrow rows or a history of white mold damage in prior seasons.



#### ONE-MONTH RISK MAP

WHAT TO DO?	
HIGH	Treat flowering fields that have a closing canopy and a history
	of white mold infections
MODERATE	Scout fields and weigh disease risk
LOW OR VERY LOW	Monitor risk forecasts



#### Risk Level

Very-low

Low

Moderate High

#### DISCI AIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Bayer, Bayer Cross and Delaro® are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.





# **Soybean Brown Spot Risk Forecast**

For period of July 10 - Aug 8, Date of Issue: July 11, 2021

### **One-Month Weather Outlook**

The next 30 days will be warmer and drier than normal for the northern parts of the Corn Belt, and generally normal for the southern parts. During the next two weeks temperatures will be cooler than normal with higher than normal precipitation for areas south of I-80. North of I-80 it will be warmer and dryer than normal except for the northeastern areas.

### **Brown Spot Risk Summary**

The risk of soybean brown spot remains moderate in the Corn Belt with growing high-risk areas.


## **Next two-week risk interpretation**

For the next two weeks the risk of soybean brown spot is moderate across the Corn Belt with high-risk areas in eastern KS, northern MO, and western IL. Scout fields in yellow, especially fields which had severe brown spot in prior seasons. Consider treatment when the disease continues to climb in the canopy.



**ONE-MONTH RISK MAP** 

WHAT TO DO?	
HIGH	Consider treatment, especially susceptible cultivars
MODERATE	Scout fields and weigh disease risk factors
LOW OR VERY LOW	Monitor risk forecasts



Very-low

Low

#### DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Bayer, Bayer Cross and Delaro® are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.



RESTRICTED

Moderate High





# Soybean Frogeye Leaf Spot Risk Forecast

For period of July 10 - Aug 8, Date of Issue: July 11, 2021

#### **One-Month Weather Outlook**

The next 30 days will be warmer and drier than normal for the northern parts of the Corn Belt, and generally normal for the southern parts. During the next two weeks temperatures will be cooler than normal with higher than normal precipitation for areas south of I-80. North of I-80 it will be warmer and dryer than normal except for the northeastern areas.

# **Frogeye Leaf Spot Risk Summary**

The risk of soybean frogeye leaf spot is moderate to high in southern areas of the Corn Belt.


## **Next two-week risk interpretation**

During the next two weeks the risk of FLS is high in parts of MO and the Delta region, and moderate in the southern Corn Belt. In areas at risk consider spraying susceptible cultivars. Scout for developing lesions on the upper leaves, especially in fields with a history of FLS infections.



**ONE-MONTH RISK MAP** 

WHAT TO DO?	
HIGH	Consider treatment, especially susceptible cultivars
MODERATE	Scout fields and weigh disease risk factors
LOW OR VERY LOW	Monitor risk forecasts



#### **DISCLAIMER**

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Bayer, Bayer Cross and Delaro® are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.

**Risk Level** 



