



DELARO[®]
Complete
 458 SC Fungicide

Soybean White Mold Risk Forecast

For period of July 31 – Aug 29, Date of Issue: August 1, 2021

One-Month Weather Outlook


Over the next 30 days temperatures will be warmer than normal north of I-80 and normal farther south. It will be dryer west of I-35, and wetter than normal in OH, IN and KY. During the next two weeks temperatures are predicted to be warmer than normal for the entire Corn Belt. Rainfall will be higher than normal east of the Mississippi River, and normal or dryer for areas west of the River.

White Mold Risk Summary

The risk of soybean white mold is moderate to high near the Great Lakes.


Next two-week risk interpretation

The conditions are moderately or highly favorable for white mold in the areas surrounding the Great Lakes during the next two weeks. In at-risk areas consider treatment if the soybeans are still flowering, especially in fields with a history of the disease.



ONE-MONTH RISK MAP

WHAT TO DO?	
HIGH	Treat flowering fields that have a closing canopy and a history of white mold infections
MODERATE	Scout fields and weigh disease risk factors
LOW OR VERY LOW	Monitor risk forecasts



Risk Level



DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Delaro[®] are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.



RESTRICTED



Soybean Brown Spot Risk Forecast

For period of July 31 – Aug 29, Date of Issue: August 1, 2021

One-Month Weather Outlook

Over the next 30 days temperatures will be warmer than normal north of I-80 and normal farther south. It will be dryer west of I-35, and wetter than normal in OH, IN and KY. During the next two weeks temperatures are predicted to be warmer than normal for the entire Corn Belt. Rainfall will be higher than normal east of the Mississippi River, and normal or dryer for areas west of the River.

Brown Spot Risk Summary

The risk of soybean brown spot slightly decreases across the Corn Belt.


Next two-week risk interpretation

During the next two weeks conditions are moderately favorable for brown spot. In at-risk areas scout fields with a history of brown spot infections in prior seasons. If brown spot keeps moving up in the canopy, consider spraying.



ONE-MONTH RISK MAP

WHAT TO DO?	
HIGH	Consider treatment, especially susceptible cultivars
MODERATE	Scout fields and weigh disease risk factors
LOW OR VERY LOW	Monitor risk forecasts



Risk Level



DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Delaro® are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.



RESTRICTED



Soybean Frogeye Leaf Spot Risk Forecast

For period of July 31 – Aug 29, Date of Issue: August 1, 2021

One-Month Weather Outlook

Over the next 30 days temperatures will be warmer than normal north of I-80 and normal farther south. It will be dryer west of I-35, and wetter than normal in OH, IN and KY. During the next two weeks temperatures are predicted to be warmer than normal for the entire Corn Belt. Rainfall will be higher than normal east of the Mississippi River, and normal or dryer for areas west of the River.

Frogeye Leaf Spot Risk Summary

The risk of disease remains moderate to high in the southern areas of the map.


Next two-week risk interpretation

For the next two weeks weather conditions are highly favorable or moderately favorable for FLS in the southern Corn Belt. In at-risk areas consider spraying susceptible cultivars that have yet to reach R5. Scout susceptible cultivars and spray if FLS is found on the upper leaves.



ONE-MONTH RISK MAP

WHAT TO DO?	
HIGH	Consider treatment, especially susceptible cultivars
MODERATE	Scout fields and weigh disease risk factors
LOW OR VERY LOW	Monitor risk forecasts



DISCLAIMER

The forecast herein is made with state of the art computer modeling technology. However, the models do not guarantee accuracy or certainty of information because of the uncertainty in climate forecast which is used in disease modeling and because of environmental variation. We specifically disclaim all warranties expressed or implied with respect to the use of this information. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Bayer, Bayer Cross and Delaro® are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2020 Bayer Group. All rights reserved.

Risk Level

